

A Mini-Review for COVID-19 issue (2020)

PAK1-blocking Therapy of Pandemic Coronaviral Infection

Hiroshi Maruta^{1} and Hong He²*

¹ PAK Research Center, Melbourne, Australia

² University of Melbourne Hospital, Melbourne, Australia

**Corresponding author: Hiroshi Maruta, Maruta20420@yahoo.co.jp*

PAK1-blocking Therapy of Pandemic Coronaviral Infection

Hiroshi Maruta^{1*} and Hong He²

¹ PAK Research Center, Melbourne, Australia

² University of Melbourne Hospital, Melbourne, Australia

****Corresponding author: Hiroshi Maruta, Maruta20420@yahoo.co.jp***

Keywords: PAK1, pathogenic kinase, viral infection, coronavirus (COVID-19), cancer

Highlights:

- 1. PAK1 is the major “pathogenic” kinase essential for infection of many viruses including COVID-19.**
- 2. Several PAK1-blockers such as propolis, melatonin, anti-malaria drugs, and ketorolac are readily available in the market.**
- 3. PAK1-blockers, interfering with the infectious process as well as promoting immune system, could be even better than traditional vaccine.**

Abstract:

PAK1 (RAC/CDC42-activated kinase 1) is the major “pathogenic” kinase whose abnormal activation causes a wide variety of diseases/disorders including cancers, inflammation, malaria and pandemic viral infection including influenza, HIV and corona virus called COVID-19. Traditionally a series of “specific” vaccines have been used for treatment of viral infection, mainly because antibiotics in general are ineffective for treatment of viral infection. However, it takes 12-18 months till the effective vaccine becomes available. Thus, as alternative instant therapeutics for COVID-19 infection, several natural and synthetic PAK1-blockers such as propolis, melatonin, **hydroxy chloroquine (HQ) and ketorolac, which are readily available in the market, are introduced in this mini-review.**

Introduction:

Like Polio, HIV, Ebola and Flu viruses, the currently pandemic virus coined COVID-19 is among RNA virus family called “corona” (Gordon et al, 2020; Maruta, 2020). Since RNA viruses need their RNA-dependent RNA polymerase (RdRP) for their replication, in theory, COVID-19 pandemic could be treated effectively by a series of RdRP inhibitors such as Remdesivir which is an ATP antagonist and originally developed in 2019 by Gilead Sciences in California for the treatment of Ebola virus (Gordon et al, 2020). However, ATP antagonists in general could potentially inhibit many other ATP-dependent enzymes such as ATPases, protein kinases, and Chaperons, and therefore could cause a number of side effects, depending on their doses, as “conventional”

anti-cancer chemos (DNA/RNA poisons). Thus, ideally, far more selective chemical drugs, which target a specific “host” enzyme essential for viral infection, but not for normal physiology of hosts, would be desirable for the COVID-19 treatment as well. Among such target enzymes is the major “pathogenic” kinase PAK1 that is essential for malaria and viral infection in general (Maruta, 2014).

Mammalian family kinases called PAKs (RAC/CDC42-activated kinases) were cloned more than 25 years ago. Among them PAK1 is the major “pathogenic” kinase whose abnormal activation is responsible for a wide variety of diseases/disorders such as cancers, inflammation, viral infection, malaria, immuno-suppression, ageing, and so on (Maruta 2014). Among PAK1-blockers, caffeic acid (CA) and its ester (caffeic acid phenethyl ester=CAPE) in a bee-product called “propolis” were the first natural ingredients that were shown to inhibit RAC, which activates directly PAK1 (Xu et al., 2005). Interestingly in 2005, an old anti-malaria drug “Chloroquine” (CQ) was also shown to suppress SARS/ coronavirus infection in cell culture with IC_{50} around $1\ \mu M$ (Vincent et al, 2005), although the precise molecular mechanism underlying its anti-viral activity remains unknown till recently. The anti-coronaviral effect of CQ and Remdesivir (IC_{50} around $1\ \mu M$ of each) was recently confirmed in vitro by a team at Chinese Academy of Sciences as well: <https://www.nature.com/articles/s41422-020-0282-0/>

In 2016, a Korean team found that the CQ up-regulates p21 (a CDK inhibitor) whose expression is suppressed by PAK1 (Oh et al, 2016; Maruta, 2014). More recently, a tumor-suppressing phosphatase called PTEN, that inactivates PAK1, was shown to suppress the coronavirus-induced LLC2-dependent fibrosis (lung inflammation) (Lu et al, 2020). Further-more, expression of LLC2 depends on the coronavirus receptor (called ACE2= Angiotensin-converting enzyme 2) -induced CK2/RAS-PAK1-RAF-AP1 signaling pathway (Chen et al, 2010, [Fig. 1](#)). These observations altogether clearly indicate the PAK1-dependency of coronaviral infection, and strongly suggest, if not proven clinically as yet, that PAK1-blockers in general could be useful for the treatment of current “pandemic” COVID-19 infection outbreak from Wohan in China since the end of 2019, which infected over [900,000](#) people world-wide, and whose death toll has reached over [44,000](#) people till now: <https://corona.help/>

In addition, PAK1 is normally responsible for the suppression of immune system in hosts (Huynh et al, 2017). Thus, like viral vaccine, these PAK1-blockers could boost the immune system for the production of antibody against this virus as well ([see Fig. 2](#)).

Natural PAK1-blockers

Traditionally a specific vaccine (based on Louis Pasteur's approach against rabies virus in 1885) is an effective cure for each viral infection. However, it takes at least 12 months (and realistically 18 months) to prepare its vaccine (according to the WHO). Until then the majority of these coronavirus victims would perish. Thus, here as "alternative" or "unconventional" therapeutics of coronaviral infection, several natural or synthetic PAK1-blockers readily available in the market are introduced.

The bee product "propolis"

Among them the bee product called "propolis" is the most popular and ancient as well. It has been used as a traditional medicine for more than 4 thousand years since the ancient Egyptian era. The father of medicine in ancient Greece, Hippocrates, was the first to coin this bee product (an alcohol-extract of beehives) "propolis" ("pro" for protection, and "polis" for beehive or city). Originally honey-bees extract something from young buds of trees such as poplar and willow and blend the extract with their saliva to make the hexagonal beehive to protect their larva from various pathogens. Thus, propolis is a "herbal" medicine prepared by bees. It is both anti-bacterial and anti-viral. It is well known to be used as a mixture of antibiotics for preparing mummies of deceased royal families to be stored under pyramids.

In modern era, propolis was recognized as an anti-cancer medicine in late 1980s by a team at Columbia University in NYC (Grunberger et al, 1988). The major anti-cancer ingredient in Egyptian or Israeli propolis turned out to be CAPE, an ester of caffeic acid (Grunberger et al, 1988) which was later known to down-regulate RAC, thereby inactivating PAK1 (Xu et al., 2005). Interestingly, however, the anti-cancer ingredients in propolis vary from one product to another, depending on where bees harvest the extract. The major anti-cancer ingredient in Brazilian green propolis is artepillin C (ARC), whereas those in subtropical propolis from Okinawa or Taiwan are polyphenols called Nymphaeols which directly inhibit PAK1 (Nguyen et al, 2017). What is common to all propolis is that they contain PAK1-blockers without any exception.

Since PAK1 is responsible not only for cancers, but also for infection with a wide variety of viruses such as influenza, HIV, pappiloma virus and SARS/coronal virus in generally, as well as immune-suppression (Maruta

2014; Huynh et al, 2017), propolis would be useful for blocking coronavirus-induced fibrosis of lungs and stimulating the immune system as well.

However, the potency of propolis varies from one product to another, depending on both chemical nature of ingredients and their content. Among propolis in the market, so far the CAPE-based New Zealand propolis called “Bio 30” (alcohol-free liquid, 25%) is the most potent (Maruta 2014). Its recommended daily dose is 1 ml (250 mg) /10 kg (body weight). Unfortunately, the cell-permeability of both caffeic acid (CA) and ARC is rather poor, mainly due to their COOH moiety. Thus, a few years ago, via Click Chemistry (CC), we managed to boost their cell-permeability by making their 1,2,3-triazolyl esters (called 15A and 15C), which are 100 and over 400 times more potent than ARC and CA, respectively (Maruta and Ahn, 2017).

Pineal hormone “melatonin”

Melatonin, a serotonin derivative from pineal glands was first recognized as an anti-melanogenic hormone by Aaron Lerner at Yale University in 1953 (Reiter and Robinson, 1995). A few years ago, we found that melanogenesis in fact depends on PAK1 (Be Tu et al, 2017). Melatonin shares a wide variety of other anti-PAK1 activities such as anti-cancer, immune stimulative, anti-infectious, anti-inflammatory, analgesic, sleepy etc. Thus, it is almost certain that melatonin, a popular sleeping pill for jet-lag treatment, could be very useful for the treatment of coronaviral infection as well.

Triptolide from thunder god vine and its water-soluble derivative

A herbal triterpene or steroid called “Triptolide “ from thunder god vine, a Chinese traditional medicine, was also found to inactivate RAC, thereby blocking PAK1 (Maruta and Ahn, 2017). Interestingly, more than a decade ago, triptolide was found to suppress virus production during dengue virus infection of human lungs by blocking PAK1 signaling pathway (Liou et al, 2008). However, its water-solubility is rather poor. Thus, several years ago, a team at University of Minnesota led by Gunda George, a German organic chemist, phosphorylated OH group at position 14 of Triptolide to boost its water-solubility over 3000 times (Patil et al, 2015). The resultant phosphatase-sensitive prodrug of triptolide called “Minnelide” is currently in clinical trials for cancers. Thus, both Triptolide and Minnelide would be potentially useful for treating coronaviral infection as well.

Ivermectin from soil bacterium (*Streptomyces avermitilis*)

Around 1975 Avermectin (a precursor of Ivermectin) was discovered from a soil bacterium by a team led by Satoshi Omura at Kitasato Institute in Tokyo, but it causes a side effect. Thus, to reduce its side effect, a Merck team led by William Campbell chemically reduced it to develop “Ivermectin” (dihydro-Avermectin), sharing the Nobel prize in 2015. It was eventually marketed by Merck into medical use in 1981. It has been used to treat many types of parasite infestations including head lice, scabies, river blindness (onchocerciasis), etc. 3 decades after its discovery, it was shown by a Russian team to suppress the growth of cancers as well, and eventually we found that inactivation of PAK1 is the major molecular mechanism underlying its anti-cancer action (Hashimoto et al, 2009). Thus, it could potentially serve as an alternative (and inexpensive) therapeutic to eradicate the PAK1-dependent coronaviral infection as well.

Artemisinin: anti-malaria from an old Chinese medical herb

In 2015 a Chinese scientist, Youyou Tu at Chinese Academy of Science, shared a Nobel prize in medicine for her discovery of anti-malaria compound called “Artemisinin” (AM). This compound was originally isolated by her “523” project team from the plant Artemisia annua, sweet wormwood, a herb employed in Chinese traditional medicine around 1972. Although the precise molecular mechanism underlying its anti-malaria and anti-viral action still remains rather unclear, the target is not the pathogens (Plasmodium falciparum or virus) themselves, but some thing in host cells, most likely PAK1 or its up-stream essential for both malaria and viral infection (Chen et al, 2009; Sicard et al, 2012), based on the following observations: (i) the AM suppresses both RAS (up-stream of PAK1) and RAF (down-stream of both RAS and PAK1) in T-cells (Maruta, 2014; Wang et al, 2007), and (ii) the dihydro derivative of AM suppresses the growth pancreatic cancer cells by up-regulation of p21 (a CDK inhibitor) whose expression is suppressed by PAK1 (Maruta, 2014; Chen et al, 2009).

Extract of Chinese (Sichuan) Pepper (Hua Jiao)

Chinese reddish peppercorns from Sichuan Province called “Hua Jiao” are among traditional spices used for the preparation of an old spicy Chinese cuisine called “Marbo-beancurd”. In 2006, we found that 70% ethanol or hot (above 45°C) water extract of Hua Jiao inhibits PAK1 with IC₅₀ around 10 µg/ml, and thereby suppressing cyclin D1 expression in both NF1-deficient triple negative breast cancer (MDA-MB-231) and MPNST cell lines in which PAK1-is abnormally activated. Just like the propolis “Bio 30”, this reddish extract (110 mg/kg, twice a week) strongly suppresses the growth of these cancer xenografts in mice (Hirokawa et al, 2006). Thus, it is most

likely that daily drinking of an inexpensive “Hua Jiao” tea (extract) could potentially contribute to both prevention and cure of COVID-19 infection, although its major PAK1-blocking ingredient has not been chemically identified as yet. For detail of health benefits from this peppercorns (promoting immune system, suppressing inflammation etc), click the following website: <https://www.organicfacts.net/sichuan-pepper.html>

FK228 (Istodax): Blocking HDAC-PAK1 pathway

A ring peptide called FK228 was isolated from a soil bacterium by a team of Fujisawa Pharm around 1995. It strongly inhibits the growth of so-called RAS cancers such as pancreatic and colon cancers which carry oncogenic mutant of Ki-RAS (Nakajima et al, 1998). A few years later it was found to inhibit directly HDAC (histone deacetylase) with IC_{50} around 1 nM (Nakajima et al, 1998). Around 2005, we found that FK228 inactivates PAK1 in several cancer cells including Tamoxiphen-resistant breast cancers and NF1-deficient MPNST (malignant peripheral nerve sheath tumor) with IC_{50} below 1 nM in cell culture, and suppresses their growth with IC_{50} below 2.5 mg /kg (i.p., twice a week) in xenografts in mice (Hirokawa et al, 2005). Around 2009, it was approved by FDA for the treatment of a rare cancer called Cutaneous T-cell Lymphoma (CTCL) etc, and sold by Celgene under the brand name “Istodax”. Thus, it is most likely useful for the therapy of COVID-19 infection as well.

Synthetic PAK1-blockers

Anti-malaria drugs: Chloroquine (CQ), and Hydroxychloroquine (HQ)

As previously described, CQ suppresses SARS/coronaviral infection in cell culture with IC_{50} around 1 μ M (Vincent et al, 2005), by upregulating p21 whose expression is suppressed by PAK1 (Oh et al, 2016, Maruta, 2014). However, due to CQ-resistance, CQ is no longer used for malaria treatment, Instead, its derivative called Hydroxychloroquine (7-chloro-4(4-(N-ethyl-N-B-hydroxyethylamino)-1-methylbutylamino)-quinoline diphosphate (HQ) has been more widely used for malaria treatment. Thus, *shortly after one of us (HM) urged both the White House and NIH to use a series of PAK1 blockers including HQ for the treatment of coronaviral infection, FDA quickly approved to use HQ or CQ for clinical trials for COVID-19 patients in NY City, the hottest-bed of COVID-19 in US. According to a French team’s clinical trial for 30-40 COVID-19 patients, the effective daily dose of HQ was found to be around 600 mg (see Fig. 3, Gautret et al, 2020).*

Ketorolac: an old pain-killer

An old pain-killer called “Toradol” is a racemic (1:1) mixture of S- and R-forms of ketorolac. Since S-form directly inhibits COX-2 which is essential for synthesis of prostaglandin, it has been used as a pain-killer. However, a few years ago, R-form was found to down-regulate RAC, thereby inactivating PAK1 (Guo et al, 2015). Thus, “Toradol” also could be used for the treatment of PAK1-dependent coronaviral infection. However, due to its COOH moiety, its cell-permeability is rather poor (with IC_{50} around 13 μ M against A549 lung cancer cell line). Thus, via Click Chemistry (CC), we have boosted its cell-permeability over 500 times (with IC_{50} around 5-24 nM against B15F10 melanoma and A549 lung cancer cells, respectively) (Nguyen et al, 2016). The resultant potent PAK1-blocker, called 1,2,3-triazolyl ester of Ketorolac (15K), suppresses both growth and metastasis of chemo-resistant human pancreatic cancer xenografts in mice with IC_{50} below 0.1 mg/kg/day, and causes no side effect even at 5 mg/kg/day (Hennig et al, 2019). Thus, 15K could be used not only for pancreatic cancer therapy, but also for therapy of infectious diseases caused by coronavirus (COVID-19) and many other deadly viruses in the future.

Vitamin D3 and its derivative (MART-10)

The most widely known pharmacological activity of Vitamin D3 is calcemic, i.e., stimulating the absorption of calcium into bone tissues. However, around late 1980s, a team in Melbourne found that Vitamin D3 can suppress the growth of cancers in mice fed with calcium-less diet (Eisman et al, 1987). However, clinically therapy of cancers with Vitamin D3 has never been successful. The main reason for its clinical failure is that Vitamin D3 is inactivated by an enzyme called CYP24 which hydroxylates at position 24 of Vitamin D3 in human body.

Thus, around a decade ago, a Japanese group led by Atsushi Kittake at Teikyo University in Tokyo developed a derivative called “MART-10” which is very resistant to CYP24 and clearly less calcemic (Flanagan et al, 2009). The “MART-10” is 1000 times more effective than Vitamin D3 in human breast and pancreatic cancers (Flanagan et al, 2009). Interestingly, a few years ago, a German group at Tuebingen University found that Vitamin D3 down-regulates RAC, thereby inactivating PAK1 and leading to depolymerization of actin filaments (Zeng et al, 2016). Independently MART-10 was also found to induce the depolymerization of actin in cancer cells (Chiang et al, 2016). Furthermore, CYP24 expression turned out to depend on the oncogenic RAS-PAK1-

NFκB/Ets signalling pathway (Dwivedi et al, 2000). Thus, it is most likely that either “MART-10” alone or a combination of Vitamin D3 and a CYP24-resistant PAK1-blocker such as propolis could be potentially useful for the treatment of coronaviral infection.

Concluding remarks

Here are introduced not only the pre-existing natural or synthetic compounds known to inhibit PAK1 directly or its up-stream, thereby potentially blocking coronaviral infection, but also their “chemical evolution” which boosts either their cell-permeability, water-solubility or metabolism-resistance over 500 times without any side effect. Furthermore, in addition to its promotion of viral infections in general, PAK1 contributes to the suppression of both B-cell and T-cell based immune systems which normally produce a series of specific antibodies against viruses (Huynh et al, 2017). Thus, PAK1-blockers could knock-out each virus with “double” punches (Fig. 2), and even in the long run could be potentially far more effective than its specific vaccine or a series of RdRP inhibitors such as Remdesivir. Lastly it should be high-lighted that the major purpose of PAK1-blockers for clinical application such as therapy of viral infection is to reduce the virus-induced abnormally activated PAK1 to the “normal” level (instead of “null” level) in patients suffering from PAK1-dependent diseases. Thus, it is very unlikely that such a cautious measure/approach would cause any serious side effect.

Conflict of Interest:

None of the authors has any conflict of interest.

References:

Be Tu PT, Nguyen BC, Tawata S et al (2017). The serum/PDGF-dependent "melanogenic" role of the minute level of the oncogenic kinase PAK1 in melanoma cells proven by the highly sensitive kinase assay. Drug Discov Ther, 10, 314-322.

Chen H, Sun B, Pan S, Jiang H, et al. (2009). Dihydroartemisinin inhibits growth of pancreatic cancer cells in vitro and in vivo. Anticancer Drugs. 20, 131-140.

Chen IY, Chang SC, Wu HY, et al (2010). Upregulation of the chemokine (C-C motif) ligand 2 (CCL2) via a severe acute respiratory syndrome coronavirus spike-ACE2 signaling pathway. *J Virol.* 84, 7703-7712.

Chiang KC, Yeh TS, Chen SC, et al. (2016). The Vitamin D Analog, MART-10, Attenuates Triple Negative Breast Cancer Cells Metastatic Potential. *Int J Mol Sci.* 17, pii: E606.

Dwivedi PP, Omdahl JL, Kola I, Hume DA, May BK. (2000). Regulation of rat cytochrome P450C24 (CYP24) gene expression. Evidence for functional cooperation of Ras-activated Ets transcription factors with the vitamin D receptor in 1,25-dihydroxyvitamin D(3)-mediated induction. *J Biol Chem.* 275, 47-55.

Eisman JA, Barkla DH, Tutton PJ. (1987). Suppression of in vivo growth of human cancer solid tumor xenografts by 1,25-dihydroxyvitamin D3. *Cancer Res.* 47, 21-25.

Flanagan JN, Zheng S, Chiang KC, et al. (2009). Evaluation of 19-nor-2alpha-(3-hydroxypropyl)-1alpha,25-dihydroxyvitamin D3 (MART-10) as a therapeutic agent for androgen-dependent prostate cancer. *Anticancer Res.* 29, 3547-3553.

Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine (HQ) and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents.* 2020 Mar 20:105949.

Gordon CJ, Tchesnokov EP, Feng JY, et al (2020). The antiviral compound remdesivir potently inhibits RNA-dependent RNA polymerase from Middle East respiratory syndrome coronavirus. *J Biol Chem.* pii: jbc.AC120.013056.

Grunberger D., Banerjee R., Eisinger K., et al. (1988). Preferential cytotoxicity on tumor cells by caffeic acid phenethyl ester isolated from propolis. *Experientia*, 44, 230-232.

Guo Y, Kenney SR Jr, Muller CY, et al. (2015). R-ketorolac Targets Cdc42 and Rac1 and Alters Ovarian Cancer Cell Behaviors Critical for Invasion and Metastasis, *Mol. Cancer Ther.* 14, 2215–2227.

Hashimoto H, Messerli SM, Sudo T, Maruta H (2009). Ivermectin inactivates the kinase PAK1 and blocks the PAK1-

dependent growth of human ovarian cancer and NF2 tumor cell lines. *Drug Discov Ther.* 3, 243-246.

Hennig R, Albawardi A, Almarzooqi S, et al. (2019). 1,2,3-Triazolyl Ester of Ketorolac (15K), a Potent PAK1-Blocker, Inhibits Both Growth and Metastasis of Human Pancreatic Cancer Orthotopic Xenografts in Mice. *Drug Discov. Ther.* 13, 248-255.

Hirokawa Y, Arnold M, Nakajima H, Zalberg J, Maruta H. (2005). Signal therapy of breast cancer xenograft in mice by the HDAC inhibitor FK228 that blocks the activation of PAK1 and abrogates the tamoxifen-resistance. *Cancer Biol. Ther.* 4, 956-960.

Hirokawa Y, Nheu T, Grimm K, et al. (2006). Sichuan pepper extracts block the PAK1/cyclin D1 pathway and the growth of NF1-deficient cancer xenograft in mice. *Cancer Biol Ther*; 5: 305-309.

Huynh N, Wang K, Yim M, et al. (2017). Depletion of p21-activated kinase 1 (PAK1) up-regulates the immune system of APC Δ 14/+ mice and inhibits intestinal tumorigenesis. *BMC Cancer.* 17, 431.

Liou JT, Chen ZY, Ho LJ, et al. (2008). Differential effects of triptolide and tetrandrine on activation of COX-2, NF- κ B, and AP-1 and virus production in dengue virus-infected human lung cells. *Eur J Pharmacol.* 589, 288-298.

Lu S, Strand KA, Mutryn MF, et al (2020). PTEN (Phosphatase and Tensin Homolog) Protects Against Ang II (Angiotensin II)-Induced Pathological Vascular Fibrosis and Remodeling-Brief Report. *Arterioscler Thromb Vasc Biol.* 40, 394-403.

Maruta H. (2014). Herbal therapeutics that block the oncogenic kinase PAK1: a practical approach towards PAK1-dependent diseases and longevity. *Phytother Res.* 28, 656-672.

Maruta H, Ahn MR. (2017). From bench (laboratory) to bed (hospital/home): How to explore effective natural and synthetic PAK1-blockers/Longevity-promoters for cancer therapy, *Eur. J. Med. Chem.* 142, 229-243.

Maruta H (2020). Tackling the Coronaviral Infection: Blocking Either the “Pathogenic” Kinase PAK1 or RNA-dependent RNA Polymerase (RdRP). *J. Infect. Dis. Ther.* 8 (2) 418.

Nakajima H, Kim YB, Terano H, Yoshida M. & Horinouchi S. (1998). FK228, a potent anti-tumor antibiotic, is a novel histone deacetylase inhibitor. *Exp Cell Res.* 241, 126-33.

Nguyen BC, Takahashi H, Uto Y, Shahinozzaman MD, Tawata S, Maruta H. (2016). 1,2,3-Triazolyl ester of Ketorolac: A "Click Chemistry"-based highly potent PAK1-blocking cancer-killer. *Eur J Med Chem.* 126, 270-276.

Nguyen BC, Yoshimura K, Kumazawa S, Tawata S, Maruta H, (2017). Frondoside A from Sea Cucumber and Nymphaeols from Okinawa Propolis: Natural anti-cancer agents that selectively inhibit PAK1 in vitro, *Drug Discov. Ther.* 11, 110-114.

Oh S, Shin JH, Jang EJ, et al. (2016). Anti-inflammatory activity of chloroquine and amodiaquine through p21-mediated suppression of T cell proliferation and Th1 cell differentiation. *Biochem Biophys Res Commun.* 474, 345-350.

Patil S, Lis LG, Schumacher RJ, et al. (2015). Phosphonooxymethyl Prodrug of Triptolide (Minnelide): Synthesis, Physicochemical Characterization, and Efficacy in Human Colon Adenocarcinoma and Ovarian Cancer Xenografts, *J. Med. Chem.* 58, 9334–9344.

Reiter R, Robinson J (1995). *Melatonin: Your Body's Natural Wonder Drug*. Bantan Books, New York.

Sicard A, Semblat JP, Doerig C, Hamelin R, et al (2011). Activation of a PAK-MEK signalling pathway in malaria parasite-infected erythrocytes. *Cell Microbiol.* 13: 836-45.

Vincent MJ, Bergeron E, Benjannet S, et al. (2005). Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology* 339, 2-12.

Wang, JX., Tang, W., Shi, LP., Wan, J. et al. (2007). Investigation of the immune-suppressive activity of artemether on T-cell activation and proliferation. *Brit. J. Pharmacol.* 150, 652-661.

Xu JW, Ikeda K, Kobayakawa A, Ikami T, et al. (2005). Downregulation of Rac1 activation by caffeic acid in aortic smooth muscle cells. *Life Sci.* 76, 2861-2872.

Zeng N, Salker MS, Zhang S, et al. (2016). $1\alpha,25$ (OH) $2D_3$ Induces Actin Depolymerization in Endometrial Carcinoma Cells by Targeting RAC1 and PAK1. *Cell Physiol Biochem*. 40, 1455-1464.

Figures:

Fig. 1. PTEN, a PAK1-blocker, interferes with coronavirus-induced PAK1-dependent signalling pathway leading to lung fibrosis.






Fig. 2. “Double” blows of PAK1-blockers against viral infection



Hydroxychloroquine (HQ) Sulfate



Effective daily dose for COVID-19: 600 mg

(Gautret P. et al , 2020)

Fig. 3.